



Turbulence and transport in fusion plasma

Nielsen, Anders Henry; Juul Rasmussen, Jens; Naulin, Volker; Madsen, Jens; Garcia, Odd-Erik

Publication date:
2008

[Link back to DTU Orbit](#)

Citation (APA):

Nielsen, A. H., Juul Rasmussen, J., Naulin, V., Madsen, J., & Garcia, O-E. (2008). *Turbulence and transport in fusion plasma*. Abstract from Danish Physical Society Annual Meeting 2008, Nyborg Strand, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Turbulence and transport in fusion plasma

By Anders H. Nielsen [1]

and

Jens Juul Rasmussen [1], Volker Naulin [1], Jens Madsen [1] and Odd-Erik Garcia [2]

[1] PLF, Risø National Laboratory for Sustainable Energy DTU, DK-4000 Roskilde

[2] Department of Physics and Technology, University of Tromsø, N-9037 Tromsø, Norway

Presentation type: talk

Turbulent transport is the dominating mechanism for plasma transport across a confining magnetic field in hot plasmas. This transport is usually referred to as anomalous transport in contrast to normal (classical/neo-classical) transport mediated by collisions. It is the limiting factor for plasma confinement in fusion plasma experiments and an important mechanism for mixing and transport in space plasmas. The turbulent fluctuations arise spontaneously due to pressure gradients perpendicular to the magnetic field. Low frequency plasma turbulence is well described by the fluid model for plasma dynamics, and it is essentially quasi two-dimensional, i.e., the turbulence is strongly anisotropic with the main dynamics in the plane perpendicular to the magnetic field. In this talk I will introduce the specific features of plasma turbulence and the associated transport/mixing. As a specific example I will discuss turbulence in the boundary region (edge plasma) and scrape-off layer (SOL) region of magnetically confined plasmas, where the relative fluctuation levels in, e.g., the plasma density approach unity.